

**WARNING: KEEP OUT OF THE REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**



Ravensdown
ANIMAL HEALTH

Zinc Oxide Plus



**Contains stabilised zinc oxide
(80% elemental zinc)**

**As an aid in the prevention of facial eczema in
cattle and sheep by oral drenching, addition to
supplementary feed or pasture spraying.**

Registered pursuant to the ACVM Act 1997, No A9412

Net weight when packed **25kg**

Ravensdown Fertiliser - 100% owned by New Zealand farmers.

**WARNING: KEEP OUT OF THE REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**



**ENVIRONMENTALLY
HAZARDOUS
SUBSTANCE, SOLID,
N.O.S. (Zinc Oxide)**

Zinc Oxide Plus

Contains 99% (990g/kg) zinc oxide (ZnO) with 1% (10g/kg) stabilising/suspending agent.

As an aid in the prevention of facial eczema in cattle and sheep by oral drenching, addition to supplementary feed or pasture spraying.

UN NUMBER: 3077
PACK GROUP III
HAZCHEM 2X

In an emergency, phone (24 hours)
0800 CHEMCALL (0800 243 622)

Read label and obtain special instructions before use.

First Aid: For advice contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

Environmental Protection: Very toxic to aquatic life and harmful to terrestrial vertebrates. Avoid release to the environment.

Disposal: Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

Emergency Response: In case of spillage wear appropriate protective clothing and prevent material from entering water ways. Sweep up without creating dust. Dispose of waste safely.

Directions for Use

1. Oral Drenching

Mix 1kg Zinc Oxide Plus with 1 litre water and stir well. This produces approximately 1.2 litres of drench. If drenching is started at least 2 weeks before the spore counts reach danger levels, the long-term drenching rate should be used. If drenching is started once spore counts are high, the crisis rate should be used.

Long-term dose rates:

Daily drenching : 3ml per 100kg live-weight

Drenching every 3 days : 13ml per 100kg live-weight

Weekly drenching : 32ml per 100kg live-weight (**weekly drenching is not suitable for lactating dairy cows**)

Crisis dose rates:

Daily drenching : 4.5ml per 100kg live-weight per day

2. Addition to Supplementary Feed

Mix properly into supplementary feed to deliver 2.5g Zinc Oxide Plus per 100kg live-weight per day.

3. Pasture Spraying

Suspend Zinc Oxide Plus in a suitable quantity of water so that pasture can be sprayed ahead of the grazing animals. Application rate (per day) is 3 - 6 grams Zinc Oxide Plus per 100kg live-weight depending on % pasture utilisation.

Recommended dose rates must be followed closely. Do not dose lactating dairy cows weekly. Zinc Oxide Plus should not be administered through the drinking water as it is insoluble in water.

Contents: 25kg net when packed

Store locked up in a clean dry covered facility.



Registered pursuant to the ACVM Act 1997,
No. A9412

See www.nzfsa.govt.nz/acvm/ for registration conditions.

ERMA Approval number HSR002054

Batch Number	
Expiry Date	

Marketed and Distributed by Ravensdown Fertiliser Co-operative Limited.
Head Office, Level 1, 32 Oxford Terrace, P O Box 1049, Christchurch,
New Zealand, Phone (+64) (3) 353-4600, Fax (+64) (3) 353-4625
Email: info@ravensdown.co.nz Website: www.ravensdown.co.nz

Ravensdown Fertiliser - 100% owned by New Zealand farmers.