

HARMFUL - KEEP OUT OF REACH OF CHILDREN

ECOTOXIC



Bait

ENDURE™

A bait for the control of
slugs and snails in crops



ACTIVE INGREDIENT: Contains 50g/kg metaldehyde in the form of a pellet. Also contains a bittering agent.

Registered To:

Helarion Industries S.A.S.
35, Rue du Corps Franc Pomies
47480 Pont Du Casse
France

NZ Agent & Importer:

Agricultural Imports Direct Ltd
Level 1, 101 Station Road,
Penrose, Auckland.
Telephone: 09 525 7236

Distributed By:

Ravensdown Fertiliser Co-operative Ltd
Level 1, 32 Oxford Terrace
Christchurch, New Zealand
Telephone: 03 353 4600
Facsimile: 03 353 4625

Batch:

D.O.M:

NET CONTENTS:

15 KG

READ LABEL CAREFULLY BEFORE USING ENDURE

This product must not be used for any purpose, or in any manner, contrary to the label unless authorized under appropriate legislation.

HAZARD CLASSIFICATION: 6.1D, 6.8B, 6.9B, 9.1C, 9.3C

TOXICITY

Harmful - May be harmful if swallowed or inhaled. May cause reproductive and liver damage from repeated oral exposure.

ECOTOXIC

Harmful to aquatic organisms. Avoid contamination of any water supply with product or empty container.

Harmful to terrestrial vertebrates. Keep away from animals, poultry, ducks and domestic pets as accidental poisoning may be fatal.

HANDLING PRECAUTIONS

STORAGE

Keep out of reach of children.

Store in accordance with NZS 8409 Management of Agrichemicals.

Store in original container tightly closed in a locked, dry, cool area away from domestic animals and poultry.

Store away from food, feedstuff, seed and fertiliser.

Stores containing 1000kg of Endure are subject to signage and secondary containment. More than 1000kg require emergency response plans.

PERSONAL PROTECTION

Avoid contact with skin and eyes. Avoid inhalation of dust.

When handling wear protective clothing such as waterproof gloves, eye protection, overalls and boots.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Do not eat, drink or smoke while using.

EQUIPMENT

Apply using well maintained and accurately calibrated equipment. Flush equipment thoroughly with clean water after use.

RECORD KEEPING

Records of use as described in NZS 8409 Management of Agrichemicals must be kept.

DISPOSAL

Ensure container is completely empty. Burn in an appropriate incinerator if circumstance, such as wind direction permit. Otherwise bury in an approved landfill. Do not use container for any other purpose. Dispose of product only by using according to label directions or dispose of safely at a managed landfill.

FIRST AID

If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If eye contact occurs, wash out immediately with water. If inhaled, move the patient to fresh air immediately. Begin artificial respiration if breathing has stopped.

SPILLAGE

In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Exclude non-essential people from the area. Sweep up spilled pellets and dispose of safely according to the label directions or at an approved landfill.

EMERGENCY RESPONSE

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call 0800 CHEMCALL (0800 243 622).

SAFETY DATA SHEET

If additional safety information is required, refer to the Safety Data Sheet. For a copy contact the customer centre on 0800 100 123.

TRANSPORT

Do not carry this product on a passenger service vehicle.

SHELF LIFE

When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

GENERAL DIRECTIONS

Endure causes the irreversible destruction of the mucus cells that are essential for movement of slugs and snails.

Best results are achieved when slugs and snails are active during mild damp weather.

In situations where pest populations are high, repeated applications of Endure may be required.

WITHHOLDING PERIOD

DO NOT apply to plant parts for human consumption or plant parts that will be grazed or feed to animals. Withhold stock from treated areas for 7 days.

PROTECTION OF ANIMALS

Store safely away from animals as accidental poisoning may be fatal. Keep poultry and pets, particularly cats and dogs away from treated areas. Endure contains a bittering agent which should deter most cats and dogs. However, some animals are inquisitive and may not be deterred. If animals are poisoned, contact a veterinarian.

APPLICATION

Apply baits when crops are most vulnerable. For arable crops this is likely to be immediately after drilling and emergence. For other crops time the application to protect sensitive growth stages. For seedlings best results may be achieved by applying 5-10 days prior to planting.

RECOMMENDATIONS FOR USE

Apply baits evenly using an appropriate applicator. Baits may be scattered on the surface or applied down the slot with seed during drilling. Ensure that any spilled baits are cleaned up, including any left from filling the applicator.

DO NOT apply to the edible parts of the plant or crops where baits maybe collected with harvested produce or in animal fodder.

DO NOT heap baits.

RECOMMENDATIONS

Under conditions of heavy rainfall and high infestation levels use the higher rates. By law, the maximum application rate (stated below) must not be exceeded.

Cereals, fodder crops and other field crops:

Broadcast: Apply 4-8kg/ha applied evenly over the soil surface or applied as a band over the area to be treated. Apply baits when crops are most vulnerable. For most crops this is likely to be at or immediately after drilling and emergence.

Use the high rates where infestation levels are high or under conditions of heavy rainfall.

The use of a granule/bait applicator is recommended.

In furrow treatment: Apply 4kg/ha and use only in situations where slugs have access to the seed through the open drilling slot e.g. direct drilling. If pest pressure is high a broadcast application may also be required to protect the emerging crop.

Vegetable crops:

Broadcast: Apply 4-8kg/ha evenly over the soil surface of the area to be treated. DO NOT apply to edible parts of crops. For short rotation crops such as lettuce, apply prior to sowing or transplanting, in periods when crops are likely to be affected by slugs and snails (early spring through to late autumn and mild winters). Applications made before crops are planted will optimise control and minimise the potential for baits to lodge on edible forage.

Use high rate where pest levels are high or under conditions of heavy rainfall.

Newly Renovated Pastures:

Broadcast: Apply 4-8kg/ha evenly over the soil surface. Use the higher rate where pest levels are high or under conditions of heavy rainfall.

The use of a granule/bait applicator is recommended.

Ornamentals:

Apply 5kg/ha or use 20g per 10 square metres. Apply evenly over the area to be treated.

WARRANTY AND LIMITATION

IMPORTANT: Read this notice before opening the container.

The registrant warrants that when packed the active ingredient of this product is as specified on this label within accepted analytical tolerances. The registrant gives only those other warranties that are required by law in relation to this product and limits its liability to the cost paid by the claimant for the product. The registrant declines all liability for damage resulting from the handling, storage or use of this product. The user bears the risk from the handling, storage or use of this product. The registrant is not liable for any use of this product which is contrary to the label directions given or any Act or Regulation.

Registered Pursuant to the ACVM Act 1997 P7818

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Approved pursuant to the HSNO Act 1996, Code: 007794

See www.erna.govt.nz for approval controls.