


266mm wide x 450mm high - PMS 1215

WARNING
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



LIME FLOUR

A calcium supplement to
aid in the prevention of
hypocalcaemia (milk fever)
in dairy cows.



ACTIVE INGREDIENT:
Ultrafine Calcium
Carbonate

NET CONTENTS:
25 KG

266mm wide x 450mm high

**WARNING: KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

LIME FLOUR

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

Handling Precautions: Causes eye irritation. Avoid contact with eyes. Wash hands thoroughly after handling.

First Aid: If eye contact occurs, rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. For advice, or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

Disposal: Preferably dispose of product by use or spreading on pasture. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Bury packaging in a suitable landfill, or if appropriate recycle.

Directions For Use

1. To aid in the prevention of hypocalcaemia (milk fever) after calving

Dose rate: 100-300 grams **Lime Flour** per cow per day, given as an oral drench, dusted onto the pasture or applied to supplementary feed. Dosing should commence as soon as possible after calving to give best results.

Drenching Instructions: To suspend, mix one part of **Lime Flour** with four parts of water. Stir the suspension frequently and flush drench lines after use to minimise settling out. If mixing **Lime Flour** and magnesium oxide (Ravensdown Magnesium Oxide Fine) together, the magnesium oxide should be suspended in water and left to stand for at least several hours before the **Lime Flour** is added.

2. To provide a calcium supplement to balance low-calcium diets

Dose rate: 50-300 grams **Lime Flour** per cow per day, mixed in with supplementary feed (eg maize silage).

Mixing Instructions: Add to the feed at a rate of 60-120 grams **Lime Flour** per kilogram of feed on a wet basis (or 20-40 grams **Lime Flour** per kilogram of feed dry matter).

Lime Flour must not be used in the last 2 months of your cows' pregnancy without veterinary advice. If you require further information on specific supplementation rates please contact Ravensdown, your veterinarian, animal health adviser or nutritionist.

ERMA Approval number HSR002521.
See www.erna.govt.nz for approval conditions.

In a transport emergency call 111; Police or Fire Brigade.
For 24 hour specialist advice in an emergency, call 0800 CHEMCALL (0800 243 622).
If additional safety information is required, refer to the Safety Data Sheet (SDS)
For a copy of the SDS phone 0800 100 123.

Close and seal bag after use.
Store locked up in a clean dry covered facility
below 30°C and away from direct sunlight.

Batch Number



Marketed and Distributed by Ravensdown Fertiliser Co-operative Limited
Level 1, 32 Oxford Terrace, Christchurch 8011, New Zealand
P O Box 1049, Christchurch 8140, New Zealand
Phone (+64) (3) 353-4600, Fax (+64) (3) 353-4625
Email: info@ravensdown.co.nz

0800 100 123
www.ravensdown.co.nz

