

WARNING KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



ravensdown

Lime Flour

A calcium supplement to aid in the prevention of hypocalcaemia (milk fever) in dairy cows.

ACTIVE INGREDIENT:

90 - 95% calcium carbonate



Registrant: Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz

BATCH:
D.O.M:

NET CONTENTS
25 KG



LIME FLOUR

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS

Causes eye irritation. Avoid contact with eyes. Wash hands thoroughly after handling.

FIRST AID

If eye contact occurs, rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. For advice, or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

DISPOSAL

Preferably dispose of product by use or spreading on pasture. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Bury packaging in a suitable landfill, or if appropriate recycle.

DIRECTIONS FOR USE

1. To aid in the prevention of hypocalcaemia (milk fever) after calving

DOSE RATE

100-300 grams Lime Flour per cow per day, given as an oral drench, dusted onto the pasture or applied to supplementary feed. Dosing should commence as soon as possible after calving to give best results.

DRENCHING INSTRUCTIONS

To suspend, mix one part of Lime Flour with four parts of water. Stir the suspension frequently and flush drench lines after use to minimise settling out. If mixing Lime Flour and magnesium oxide (Ravensdown Magnesium Oxide Drenching) together, the magnesium oxide should be suspended in water and left to stand for at least several hours before the Lime Flour is added.

2. To provide a calcium supplement to balance low-calcium diets

DOSE RATE

50-300 grams Lime Flour per cow per day, mixed in with supplementary feed (eg maize silage).

Mixing Instructions: Add to the feed at a rate of 60-120 grams Lime Flour per kilogram of feed on a wet basis (or 20-40 grams Lime Flour per kilogram of feed dry matter).

Lime Flour must not be used in the last 1-2 months of your cows' pregnancy without veterinary advice. If you require further information on specific supplementation rates please contact Ravensdown, your veterinarian, animal health adviser or nutritionist.

WITHHOLDING PERIOD

Meat withholding: NIL

Milk withholding: NIL

EPA APPROVAL NUMBER HSR002521

See www.epa.govt.nz for approval conditions.

In a transport emergency call 111; Police or Fire Brigade. For 24 hour specialist advice in an emergency, call 0800 CHEMCALL (0800 243 622). If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.

Close and seal bag after use. Store locked up in a clean dry covered facility below 30°C and away from direct sunlight.

PHONE 0800 100 123
24 Hours - 7 Days