

**WARNING – KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**



ravensdown

Magnesium Oxide Drenching

A magnesium supplement to aid in
the prevention of hypomagnesaemia
in cattle

ACTIVE INGREDIENT:

325 mesh drenching-grade,
90% magnesium oxide (54%
elemental magnesium).



Marketed & distributed by Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600

ravensdown.co.nz

BATCH:

**NET CONTENTS
20 KG**

**WARNING – KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

ravensdown 

MAGNESIUM OXIDE DRENCHING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS

Causes eye irritation and may cause damage to respiratory system. Avoid contact with eyes and avoid breathing dust. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling.

FIRST AID

If eye contact occurs, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. For advice, or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

DISPOSAL

Preferably dispose of product by use or spreading on pasture. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Bury packaging in a suitable landfill, or if appropriate recycle.

DIRECTIONS FOR USE

Supplement at times such as in early spring when there are low magnesium levels, and gradually reduce daily allowance once risk periods are past. It is generally recommended to supplement a minimum of 10-15g elemental magnesium per cow per day, depending on animals' level of magnesium deficiency.

1. Oral Drenching (post-calving)

Mix thoroughly with water and during drenching shake regularly to maintain suspension. Dose with 20- 30g/cow/day to provide 10-15g of elemental magnesium per day.

2. Pasture Dusting

Best results are achieved when applied to damp pasture. Dusted pasture can tolerate light rain but re-dust if heavy rain occurs within 2-3 days of application. Apply daily to pasture about to be grazed at 50-70g/cow/day.

3. Addition to Supplementary Feed

Mix with water and apply evenly over feed at a rate of 40-60g/cow/day.

EPA APPROVAL NUMBER HSR006644

See www.epa.govt.nz for approval conditions.

In a transport emergency call 111; Police or Fire Brigade. For 24 hour specialist advice in an emergency, call 0800 CHEMCALL (0800 243 622). If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.

Close and seal bag after use. Store locked up in a clean dry covered facility below 30°C and away from direct sunlight.

PHONE 0800 100 123