

**WARNING – KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

  
**ravensdown**

# Zinc Oxide Plus<sup>TM</sup>

An aid in the prevention of facial  
eczema in cattle and sheep by oral  
drenching, addition to supplementary  
feed or pasture spraying

---

**ACTIVE INGREDIENT:**

Stabilised zinc oxide (80% elemental zinc).

---



**Registrant: Ravensdown Ltd**  
292 Main South Road  
Hornby, Christchurch 8042  
New Zealand  
Phone: 03 353 4600  
[ravensdown.co.nz](http://ravensdown.co.nz)

**NET CONTENTS**  
**25 KG**

**WARNING – KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

**ravensdown** 

# ZINC OXIDE PLUS™ | READ LABEL CAREFULLY BEFORE USING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

## FIRST AID

For medical advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

## ENVIRONMENTAL PROTECTION

Very toxic to aquatic life and harmful to terrestrial vertebrates. Avoid release to the environment.

## DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

## EMERGENCY RESPONSE

In case of spillage wear appropriate protective clothing and prevent material from entering water ways. Sweep up without creating dust. Dispose of waste safely.

## DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

## ORAL DRENCHING

Mix 1kg **ZINC OXIDE PLUS** with 1 litre of water and stir well. This produces approximately 1.2 litres of drench. If drenching is started at least 2 weeks before the spore counts reach danger levels, the long-term drenching rate should be used. If drenching is started once spore counts are high, the crisis rate should be used.

### Long-term dose rates:

Daily drenching: 3ml per 100kg live-weight  
Drenching every 3 days: 13ml per 100kg live-weight  
Weekly drenching: 32ml per 100kg live-weight (weekly drenching is not suitable for lactating cows)

### Crisis dose rates:

Daily drenching: 4.5ml per 100kg live-weight per day

## ADDITION TO SUPPLEMENTARY FEED

Mix properly into supplementary feed to deliver 2.5g **ZINC OXIDE PLUS** per 100kg live-weight per day.

## PASTURE SPRAYING

Suspend **ZINC OXIDE PLUS** in a suitable quantity of water so that pasture can be sprayed ahead of grazing animals. Application rate (per day) is 3-6 grams of **ZINC OXIDE PLUS** per 100kg live-weight depending on % pasture utilisation.

Recommended dose rates must be followed closely. Do not dose lactating dairy cows weekly. **ZINC OXIDE PLUS** should not be administered through the drinking water as it is insoluble in water.

**ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S.** (Zinc Oxide) UN Number 3077, Pack Group III, Hazchem 2Z

Registered pursuant to the ACVM Act 1997, No.A9412. See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions. EPA Approval number: HSR002054. See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions.

In a transport emergency call 111; Police or Fire Brigade. For 24 hour specialist advice in an emergency, call 0800 CHEMCALL (0800 243 622). If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.

Close and seal bag after use. Store locked up in a clean dry covered facility below 30°C and away from direct sunlight.

BATCH:

EXP:

**PHONE 0800 100 123**

