


**CAUTION** – DO NOT ALLOW DOGS, HORSES, OR OTHER EQUIDS ACCESS TO FEEDS CONTAINING BOVATEC® 20CC AS INGESTION BY THESE SPECIES COULD BE FATAL.

**ravensdown**

# RavCalf 14

## **14% protein calf pellets with Bovatec® 20CC**

- Highly palatable
- Balanced post-weaning feed
- High energy, grain based feed to support calf growth after weaning
- Contains Bovatec® 20CC as an aid in the control of coccidiosis



**Distributor: Ravensdown Ltd**  
292 Main South Road  
Hornby, Christchurch 8042  
New Zealand  
Phone: 03 353 4600  
[ravensdown.co.nz](http://ravensdown.co.nz)

**NET CONTENTS**  
**25 KG**

**CAUTION – DO NOT ALLOW DOGS, HORSES, OR OTHER EQUIDS ACCESS TO FEEDS CONTAINING BOVATEC® 20CC AS INGESTION BY THESE SPECIES COULD BE FATAL.**

**ravensdown** 

## **RAVCALF 14** | READ LABEL CAREFULLY BEFORE USING

**High energy grain based feed to support growth and development of calves that have been weaned off milk and have transitioned to a grass based diet. Contains 100mg per kg lasalocid (as 500mg/kg Bovatec® 20CC, A9679) as an aid in the control of coccidiosis caused by Eimeria species.**

### **FEEDING RECOMMENDATIONS**

Designed to be fed as a supplement to grass once calves are weaned off CMR or milk. Transition slowly from RavCalf 20 up to a maximum of 3kg per head per day. Feed 1kg of **RAVCALF 14** per 100kg calf to provide 1mg/kg lasalocid per day.

Continue to feed until target weights are reached.

Forage should always be fed when feeding pelletised calf feed. Ensure calves have unrestricted access to fresh and clean water at all times.

### **TYPICAL ANALYSIS (Dry Matter basis)**

Minimum crude protein	14%
Maximum crude fat	5%
Maximum crude fibre	10%
Maximum salt	1%

### **INGREDIENTS SELECTED FROM:**

Cereal grains and by-products, oil seed meals, molasses, vegetable fats and oils, bentonite, dicalcium phosphate, limestone, salt, trace elements, vitamins, flavouring and Bovatec® 20CC (coccidiostat).

### **BOVATEC® 20CC**

#### **FOR ANIMAL TREATMENT ONLY.**

**Contains lasalocid sodium at the rate of 100mg/kg feed.**

**Contraindications:** Care must be exercised when feeding concurrently with other antimicrobials. Do not feed with other ionophores, e.g. monensin capsules, liquid or premix. Do not exceed recommended dose rates. Not to be used for single dose treatments.

### **WITHHOLDING PERIODS**

Meat: Nil

Milk: Nil

Bovatec® 20CC is registered pursuant to the ACVM Act 1997, No.A09679. See [www.mpi.govt.nz](http://www.mpi.govt.nz) for registration details.

Store in a cool dry place out of direct sunlight.

**PHONE 0800 100 123**