

WARNING – KEEP OUT OF REACH OF CHILDREN.
FOR ANIMAL TREATMENT ONLY


ravensdown

Lift[®] B₁₂ Plain

For the treatment of cobalt deficiency and vitamin B₁₂ responsive conditions in cattle and sheep



ACTIVE INGREDIENT:

Contains 2mg/ml hydroxocobalamin (long acting vitamin B₁₂).

NET CONTENTS
500 ML

Store in a cool dry place below 30°C (room temperature) in the carton to protect product from light and keep locked up in a safe place.

Registered pursuant to the ACVM Act 1997, No.A9783.
See www.foodsafety.govt.nz for registration conditions.

Registrant: Ravensdown Ltd

292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600

ravensdown.co.nz

BATCH:

EXP:



460605 V01

PHONE 0800 100 123

ravensdown 

LIFT[®] B₁₂ PLAIN

READ LABEL CAREFULLY BEFORE USING

Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

Vitamin B₁₂ deficiencies: Before treatment, establish vitamin B₁₂ status of stock and/or pasture cobalt levels. Cobalt deficiency in soil and/or pasture leads to a lack of vitamin B₁₂ in body tissues of stock. Vitamin B₁₂ deficiency is mostly a problem in very young stock. Poor growth rates and ill thrift may be seen in affected animals. Treatment of vitamin B₁₂ deficiency is likely to last only 4 - 6 weeks. Treatment of pregnant animals should benefit newborn animals particularly when treated in the latter stages of pregnancy.

Supplementation of young ruminants should begin at or before weaning. The need for, and frequency of, dosing is related to the severity of deficiency and the age of the animal. Seek veterinary advice if poor stock condition persists after treatment.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Administer by subcutaneous or intramuscular injection, into the anterior half of the neck.

DOSE RATE

Cattle: 2-6ml

Sheep: 0.5-1ml

WITHHOLDING PERIODS

Nil

WARNING - KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY

ravensdown

Lift® B₁₂ Plain

For the treatment of cobalt deficiency and vitamin B₁₂ responsive conditions in cattle and sheep



ACTIVE INGREDIENT:

Contains 2mg/ml hydroxocobalamin (long acting vitamin B₁₂).



Registrant: Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz

BATCH:
EXP:

NET CONTENTS
500 ML

ravensdown

Lift® B₁₂ Plain

ravensdown

LIFT® B₁₂ PLAIN

**READ LABEL CAREFULLY
BEFORE USING**

Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

Store in a cool dry place below 30°C (room temperature) in the carton to protect product from light and keep locked up in a safe place.

Registered pursuant to the ACVM Act 1997, No.A9783.
See www.foodsafety.govt.nz for registration conditions.

PHONE 0800 100 123

500 ML
NET CONTENTS

Registrant: Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz



ACTIVE INGREDIENT:
Contains 2mg/ml hydroxocobalamin (long acting vitamin B₁₂).



For the treatment of cobalt deficiency and vitamin B₁₂ responsive conditions in cattle and sheep

Lift® B₁₂ Plain

ravensdown

WARNING - KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY

Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

**READ LABEL CAREFULLY
BEFORE USING**

LIFT® B₁₂ PLAIN

ravensdown

Vitamin B₁₂ deficiencies: Before treatment, establish vitamin B₁₂ status of stock and/or pasture cobalt levels. Cobalt deficiency in soil and/or pasture leads to a lack of vitamin B₁₂ in body tissues of stock. Vitamin B₁₂ deficiency is mostly a problem in very young stock. Poor growth rates and ill thrift may be seen in affected animals. Treatment of vitamin B₁₂ deficiency is likely to last only 4 - 6 weeks. Treatment of pregnant animals should benefit newborn animals particularly when treated in the latter stages of pregnancy. Supplementations should begin at or before weaning. The need for, and frequency of, dosing is related to the severity of deficiency and the age of the animal. Seek veterinary advice if poor stock condition persists after treatment.

DIRECTIONS FOR USE
By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DOSE RATE
Administer by subcutaneous or intramuscular injection, into the anterior half of the neck.

| | |
|----------------------------|----------------|
| Cattle: 2-6ml | Sheep: 0.5-1ml |
| WITHHOLDING PERIODS | |
| NIL | |

460610 V01