

WARNING – KEEP OUT OF REACH OF CHILDREN.
FOR ANIMAL TREATMENT ONLY


ravensdown

Lift[®] B₁₂ Selenised

For the treatment and prevention of vitamin B₁₂ and selenium responsive conditions in cattle and sheep



ACTIVE INGREDIENTS:

Contains 2mg/ml hydroxocobalamin (long acting vitamin B₁₂) 4mg/ml selenium (as sodium selenate anhydrous).

NET CONTENTS
500 ML

Harmful to aquatic life with long-lasting effects. Avoid release to the environment. Store in a cool dry place below 30°C (room temperature) in the carton to protect product from light and keep locked up in a safe place.

EPA approval number HSR002387. See www.epa.govt.nz for approval conditions.
Registered pursuant to the ACVM Act 1997, No.A9784.
See www.foodsafety.govt.nz for registration conditions.

Registrant: Ravensdown Ltd

292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600

ravensdown.co.nz

BATCH:

EXP:



460616 V01

PHONE **0800 100 123**

ravensdown 

LIFT[®] B₁₂ SELENISED

READ LABEL CAREFULLY BEFORE USING

Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

Vitamin B₁₂ and Selenium deficiencies: Before treatment, establish selenium and vitamin B₁₂ status of stock and/or pasture cobalt levels. Do not administer LIFT[®] B₁₂ SELENISED unless deficiencies of both cobalt and selenium exists. Cobalt deficiency in soil and or pasture leads to lack of vitamin B₁₂ in body tissues of stock. Vitamin B₁₂ deficiency is mostly a problem in very young stock. Poor growth rates and ill thrift may be seen in affected animals. Treatment of Vitamin B₁₂ deficiency is likely to last only 4 - 6 weeks. Re-dose with non-selenised vitamin B₁₂ after this time if required. Treatment of pregnant animals should benefit newborn animals particularly when treated in the latter stages of pregnancy. Supplementation of young ruminants should begin at or before weaning, the need for, and the frequency of, dosing is related to the severity of deficiency and the age of the animal. Seek veterinary advice if poor stock condition persists after treatment.

Contraindications:

Do not use unless a selenium deficiency has been diagnosed. Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. As selenium is toxic above the recommended rates, do not exceed stated dose, or dose more frequently than once every 3 months.

DIRECTIONS FOR USE

By law the user must take due care, seeking expert advice if necessary, to avoid unnecessary pain or distress in the treated animal when using the product in a manner other than as recommended on the label. Administer by subcutaneous or intramuscular injection, into the anterior half of the neck.

DOSE RATE

Calves: 2-3ml	Adult cattle: 4-6ml	Lambs: 0.5ml	Adult sheep: 1-1.5ml
---------------	---------------------	--------------	----------------------

WITHHOLDING PERIODS

Nil

WARNING - KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY

ravensdown

Lift® B₁₂ Selenised

For the treatment and prevention of vitamin B₁₂ and selenium responsive conditions in cattle and sheep



ACTIVE INGREDIENTS:

Contains 2mg/ml hydroxocobalamin (long acting vitamin B₁₂)
4mg/ml selenium (as sodium selenate anhydrous).



Registrant: Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz

BATCH:
EXP:

NET CONTENTS
500 ML

ravensdown

Lift® B₁₂ Selenised

ravensdown

LIFT® B₁₂ SELENISED

**READ LABEL CAREFULLY
BEFORE USING**

**Read label and obtain special instructions
before use. Do not handle until all safety
precautions have been read and understood.**

Harmful to aquatic life with long-lasting effects. Avoid release to the environment. Store in a cool dry place below 30°C (room temperature) in the carton to protect product from light and keep locked up in a safe place.

EPA approval number HSR002387. See www.epa.govt.nz for approval conditions. Registered pursuant to the ACVM Act 1997, No.A9784. See www.foodsafety.govt.nz for registration conditions.

PHONE 0800 100 123

NET CONTENTS
500 ML

Registrant: Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz



ACTIVE INGREDIENTS:
Contains 2mg/ml hydroxocobalamin (long acting vitamin B₁₂)
4mg/ml selenium (as sodium selenate anhydrous).

For the treatment and prevention of vitamin B₁₂ and selenium responsive conditions in cattle and sheep

Lift® B₁₂ Selenised

ravensdown

WARNING - KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY

ravensdown
LIFT® B₁₂ SELENISED
READ LABEL CAREFULLY
BEFORE USING
Read label and obtain special instructions
before use. Do not handle until all safety
precautions have been read and understood.

Vitamin B₁₂ and Selenium deficiencies: Before treatment, establish selenium and vitamin B₁₂ status of stock and/or pasture cobalt and selenium levels. Do not administer LIFT® B₁₂ SELENISED unless deficiencies of both cobalt and selenium exist. Cobalt deficiency in soil and/or pasture leads to lack of vitamin B₁₂ in body tissues of stock. Vitamin B₁₂ deficiency is mostly a problem in very young stock. Poor growth rates and ill thrift may be seen in affected animals. Treatment of vitamin B₁₂ deficiency is likely to last only 4 - 6 weeks. Re-dose with non-selenised vitamin B₁₂ after this time if required. Treatment of pregnant animals should benefit newborn animals particularly when treated in the latter stages of pregnancy. Supplementation of young ruminants should begin at or before weaning, the need for, and the frequency of, dosing is related to the severity of deficiency and the age of the animal. Seek veterinary advice if poor stock condition persists after treatment.

Contraindications: Do not use unless a selenium deficiency has been diagnosed. Do not use at the same time as any other selenised fertiliser, pill or product without consulting a veterinarian. As selenium is toxic above the recommended rates, do not exceed stated dose, or dose more frequently than once every 3 months.

460621 V02

DIRECTIONS FOR USE
By law the user must take due care, seeking expert advice if necessary, to avoid unnecessary pain or distress in the treated animal when using the product in a manner other than as recommended on the label.
Administer by subcutaneous or intramuscular injection, into the anterior half of the neck.

DOSE RATE	
Calves: 2-3ml	Adult cattle: 4-6ml
Lambs: 0.5ml	Adult sheep: 1-1.5ml

WITHHOLDING PERIODS

NIL	
-----	--