

WARNING – KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

**ravensdown**


Moximax[®]

Sheep

A broad spectrum sheep drench for the control of susceptible gastrointestinal roundworms and lungworms in sheep and lambs including benzimidazole, levamisole and morantel resistant strains

ACTIVE INGREDIENTS:

1.0g/litre moxidectin, 1.0g/litre iodine,
0.2g/litre cobalt, 2.0g/litre copper,
0.6g/litre zinc, 0.5g/litre selenium.

**Registrant: Ravensdown Ltd**

292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600

ravensdown.co.nz

BATCH:

EXP:

NET CONTENTS
20 Litres

MOXIMAX[®] SHEEP | READ LABEL CAREFULLY BEFORE USING

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

ECOTOXICITY

Very toxic to aquatic organisms. Avoid contamination of any water supply. **Harmful to the soil environment.** Harmful to terrestrial invertebrates. Avoid release into the environment.

HANDLING PRECAUTIONS

Repeated exposure may cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause harm to breastfed children. Avoid contact during pregnancy/while nursing. Avoid breathing vapours, wear protective gloves and use personal protective equipment as required. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace and should be washed before reuse.

DISPOSAL

Preferably dispose of product by use. Otherwise treat the product so that it is no longer toxic and dispose of product and packaging at an approved landfill or other approved facility. Triple (or preferably pressure) rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate, recycle.

FIRST AID

If swallowed rinse mouth and do NOT induce vomiting. If skin contact occurs wash with plenty of soap and running water. For advice or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

EMERGENCY RESPONSE

In case of spillage wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely.

EFFICACY SPECTRUM

MOXIMAX SHEEP contains 1g/L moxidectin, 0.5g/L selenium, 0.2g/L cobalt, 2g/L copper, 0.6g/L zinc and 1g/L iodine. For the treatment of the following susceptible immature and mature (includes inhibited L4) gastrointestinal roundworms and lungworms, including benzimidazole, levamisole and morantel resistant strains: *Haemonchus spp.*, *Ostertagia spp.*, *Cooperia spp.*, *Trichostrongylus spp.*, *Oesophagostomum spp.* (adults only), *Nematodirus spp.* (adults only), *Dictyocaulus spp.*, *Chabertia ovina*, *Strongyloides papillosus* (adults only), *Trichuris ovis* (adults only).

PERSISTENT ACTIVITY

When used at the recommended dose rate **MOXIMAX SHEEP** will prevent reinfection by *Haemonchus contortus* for up to 28 days and *Ostertagia circumcincta* for up to 21 days following the administration of a single oral dose.

RESISTANCE MANAGEMENT

Resistance can develop to any anthelmintic. Ask your local animal health advisor or veterinarian for recommended parasite management practices for your area to reduce the development of resistance. It is advised that a resistance test be carried out regularly when using any parasite treatment.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Do not use at the same time as any other selenised fertiliser, prill or product without consulting your animal health advisor. Do not exceed the stated dose, or dose more frequently than 3-weekly. Caution: In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (death) occurring following copper supplementation in this species.

Shake well before use. **MOXIMAX SHEEP** is a ready to use oral anthelmintic that can be dosed through standard drenching equipment. It is important to give the correct dose. Therefore, before treatment, weigh a representative sample of sheep to determine the weight of the heaviest animal. Where there is a large variation in live-weights, drafting into two or more lines may be appropriate before weighing and treating, to avoid under and over dosing. In this situation it is important to choose the dose rate based on the label recommendation for each weight band. Use correct drenching techniques and check accuracy of drenching equipment regularly. Wash out drench gun thoroughly with water before and after use.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MEAT

Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of cessation of the last treatment.

MILK

Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

DOSE RATE

1ml per 5kg live-weight orally (0.2mg/kg moxidectin).

DOSE VOLUME TABLE

Live-weight (kg)	Dose (ml)	20L treats (head)
11-15	3	6664
16-20	4	5000
21-25	5	4000
26-30	6	3333
31-35	7	2857
36-40	8	2500
41-45	9	2222
46-50	10	2000
51-55	11	1818
56-60	12	1666
61-65	13	1538
66-70	14	1428
71-75	15	1333

Sheep heavier than 75kg should be dosed at 1ml per 5kg live-weight.



Agrecovery is available on plastic containers up to 60L and for plastic or steel large drums from 61L - 1000L.

PHONE **0800 100 123**



Registered pursuant to the ACVM Act 1997, No.A10724. See www.foodsafety.govt.nz for registration conditions. Approved pursuant to the HSNO Act 1996, Code: HSRO01900. See www.epa.govt.nz for approval controls.

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (contains moxidectin)
UN Number 3082, Pack Group III, Hazchem 3Z

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call **0800 CHEMICAL (0800 243 622)**. If additional safety information is required, refer to the Safety Data Sheet. For a copy, contact the customer centre on 0800 100 123.

Store below 25°C, store in original container, tightly closed away from direct sunlight, do not freeze.