

**WARNING** – KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY - SHAKE WELL BEFORE USE

  
**ravensdown**

# Deltamethrin

## Pour On

For the prevention of  
nuisance fly, control of biting  
lice and sucking lice.

---

**ACTIVE INGREDIENT:**

15g/L Deltamethrin

---

**FLY AND LICE POUR ON  
FOR CATTLE**



**Registered to: Nexan  
Corporation Ltd**  
PO Box 217234  
Botany Junction  
Auckland, New Zealand

**Distributor: Ravensdown Ltd**  
292 Main South Road  
Hornby, Christchurch 8042  
New Zealand  
Phone: 03 353 4600

[ravensdown.co.nz](http://ravensdown.co.nz)

**NET CONTENTS  
5 Litres**

ravensdown 

## DELTAMETHRIN POUR ON | READ LABEL CAREFULLY BEFORE USING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

### WARNING: DANGEROUS TO THE ENVIRONMENT

HAZARD CLASSIFICATION: 6.3B, 6.4A, 6.5B, 6.9B, 9.1A, 9.2C, 9.4A

### SAFETY PRECAUTIONS AND DIRECTIONS

May cause eye or mild skin irritation, an allergic skin reaction and damage to nervous system from repeated exposure at high doses. Avoid skin contact, wear protective clothing and gloves. Wash hands after handling. Do not breath in vapours.

### FIRST AID

If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If eye contact occurs, hold eyes open and rinse cautiously with water for several minutes. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

### ENVIRONMENTAL PROTECTION

Harmful to aquatic organisms. Avoid contamination of any water supply with product or empty container. Avoid release to the environment.

### DISPOSAL

Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Avoid contamination of any water source. Triple rinse container immediately after emptying and pour rinsate onto waste ground. Do NOT use container for any other purpose.

### EMERGENCY RESPONSE

In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely.

### DIRECTIONS FOR USE

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.** SHAKE WELL. Apply through a suitable clean application gun with a T-bar nozzle. **Must be applied in a wide (5 to 10cm) continuous strip from between the ears to the tail, down the midline, to ensure even dispersion of product down both sides of the animal.** Dose according to the heaviest animal, see below dose rate table. Clean equipment thoroughly after use.

### DOSE RATE: LICE AND NUISANCE FLY - 1ML PER 20KG

LIVEWEIGHT	DOSE	DOSES PER 5L PACK
Up to 100kg	5ml	1000
101 - 200kg	10ml	500
201 - 300kg	15ml	333
301 - 400kg	20ml	250
401 - 500kg	25ml	200
501 - 600kg	30ml	166
Cattle over 600kg	5ml per 100kg liveweight	

BATCH:

EXP:

Agrecovery is available on plastic containers up to 60L and for plastic or steel large drums from 61L - 1000L.



PHONE 0800 100 123

### WITHHOLDING PERIODS

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food. Notice: Maximum Residue Levels for Agricultural Compounds.**

**MEAT:** Cattle producing meat or offal for human consumption must not be sold for slaughter either during or within 28 days of the last treatment.

**MILK:** Nil

**BOBBY CALVES:** No withholding period is required prior to slaughter for calves born to cows treated in late pregnancy and then allowed to suckle those cows.

### GENERAL INFORMATION


Ravensdown DELTAMETHRIN Pour On is a water based synthetic pyrethroid insecticide for control of biting and sucking lice (*Bovicola bovis*, *Linognathus vituli*, *Haematopinus eurysternus* and *Solenopotes capillatus*) on cattle of any age.

It also controls nuisance fly (*Stomoxys calcitrans*) for a period of up to four weeks. The period of protection Ravensdown DELTAMETHRIN Pour On will achieve is influenced by a number of factors such as the level of fly challenge, environmental and management factors. Therefore it is imperative that Ravensdown DELTAMETHRIN Pour On is considered part of an overall nuisance fly management plan.

Maximum lice kill is achieved up to six weeks after application, though relief of symptoms is rapid. DO NOT apply Ravensdown DELTAMETHRIN Pour On to wet cattle. Do not use if rainfall is pending.

**Ravensdown DELTAMETHRIN Pour On is a non-systemic. It controls lice by dispersing externally around the body of the animal; therefore accurate application is critical.**

**Precautions:** In rare cases some cows are irritated by synthetic pyrethroids for 48 to 72 hours after application. This tends to be on a herd basis and repeatable so do not use on cows previously known to be irritated by synthetic pyrethroids.



Registered pursuant to the HSNO Act 1996, HSRO02226. See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions. Registered pursuant to the ACVM Act 1997, No. A11646. See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Store at or below 25°C, avoid temperatures below 0°C and above 35°C. Protect from light.

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call **0800 CHEMICAL (0800 243 622)**. If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.