

**WARNING** - KEEP OUT OF REACH OF CHILDREN - FOR ANIMAL TREATMENT ONLY

  
**ravensdown**

# Flumethrin

For the control of ticks on  
cattle and as an aid in the  
treatment of ticks on deer

---

**ACTIVE INGREDIENT:**

10g/L Flumethrin

---



**POUR ON TICKICIDE  
FOR CATTLE AND DEER**



**Registered to: Nexan  
Corporation Ltd**  
PO Box 217234  
Botany Junction  
Auckland, New Zealand

**Distributor: Ravensdown Ltd**  
292 Main South Road  
Hornby, Christchurch 8042  
New Zealand  
Phone: 03 353 4600

[ravensdown.co.nz](http://ravensdown.co.nz)

**NET CONTENTS  
5 Litres**



# FLUMETHRIN | READ LABEL CAREFULLY BEFORE USING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

## HANDLING PRECAUTIONS

Ravensdown FLUMETHRIN may cause a temporary skin irritation in some individuals if applied to the skin. Avoid contact with hands and skin. Wear protective clothing and gloves when handling. Do not breathe vapour.

## FIRST AID

If on skin wash off with soap and water. If you feel unwell get medical advice. Call 0800 POISON (0800 764 766) or a doctor immediately.

## ENVIRONMENTAL PROTECTION

Harmful to aquatic life with long lasting effects. Avoid contamination of any water supply with product or empty container. Harmful to terrestrial vertebrates. Avoid release to the environment.

## DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/facility. Triple rinse empty container. Spread rinsate onto waste ground or pasture. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose.

## EMERGENCY RESPONSE

For specialist advice in an emergency call 111.

## GENERAL USE

Ravensdown FLUMETHRIN spreads through the oils of skin and hair coat of all parts of the animal's body. Behaviour such as grooming, tail swishing and scratching, aid the movement of active ingredient to the extremities of the body, which are often affected by ticks.

Ravensdown FLUMETHRIN is active against all 3 stages of the New Zealand cattle tick (*Haemaphysalis longicornis*) lifecycle. It also causes infertility of ticks which survive treatment, by inhibiting egg laying or rendering eggs sterile. This action helps reduce contamination of pasture.

## DIRECTIONS FOR USE

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.**

Apply Ravensdown FLUMETHRIN evenly along the midline of the back from the front of the shoulders to the tail head. Use a plastic measuring cup or automatic drenching gun to apply. Control of ticks is not affected by rain falling within a few hours of treatment. Retreatment should be undertaken when ticks start to reappear. For optimum control of deer, breeding hinds should be treated every three weeks from August to fawning. This program will reduce the number of nymphs reaching the adult stage, which infest fawn during November/December. Treatment of hinds can be continued after fawning if this is practical. Other stock should be treated every three weeks from August, or as required.

## ADVERSE EFFECTS

Signs of mild irritation may be observed for 24 to 48 hours after treatment. Hair loss at the site of application may occur on deer which are shedding their winter coat at the time of treatment. This effect is temporary and can be minimised by spreading the dose over the entire backline. Growth of summer coat is not affected.

## CONDITION OF SALE

The manufacturer of this animal remedy extends/grants to the purchaser a warranty that this animal remedy is reasonably fit the purposes for which its use is recommended, provides that the purchaser uses the remedy only for the purpose for which it is recommended and strictly in accordance with the directions on this container.

## DOSE RATE

CATTLE (Liveweight)		DEER	
WEIGHT RANGE	DOSE VOLUME	AGE RANGE	DOSE VOLUME
Up to 200kg	20ml	Up to 6 months	10ml
200 - 300kg	30ml		
301 - 400kg	40ml	Yearlings and adults	20ml
Over 400kg	50ml		

## WITHHOLDING PERIODS

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

**MEAT:** Nil.

**MILK:** Nil

There is **NO WITHHOLDING PERIOD** for meat and milk from treated animals

Registered pursuant to the HSNO Act 1996, HSRO01801. See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions. Registered pursuant to the ACVM Act 1997, No.A11502. See [www.foodsafety.govt.nz/acvm/](http://www.foodsafety.govt.nz/acvm/) for registration conditions.

Store below 25°C (room temperature). Do not freeze.

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call **0800 CHEMICAL (0800 243 622)**. If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.

BATCH:

EXP:

Agrecovery is available on plastic containers up to 60L and for plastic or steel large drums from 61L - 1000L.

PHONE **0800 100 123**

 **AGRECOVERY**  
[www.agrecovery.co.nz](http://www.agrecovery.co.nz)