

WARNING – KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY


ravensdown

MoximaxTM

Pour On

Long lasting endectocide for cattle and deer. For the treatment and control of internal and external parasites of cattle (including lactating dairy cattle) and for the treatment and control of lungworm and roundworms of deer.

Improves milk production in lactating dairy cows and increases weight gains in growing beef and dairy cattle exposed to parasite challenge.

ACTIVE INGREDIENT:

5g/L MOXIDECTIN
Contains aromatic
hydrocarbons 150g/L.



Registered to: Polygon (NZ) Ltd
PO Box 25179
Auckland 1740
New Zealand

Marketed & distributed by:
Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz

NET CONTENTS
5 Litres

ravensdown 

MOXIMAX™ POUR ON | READ LABEL CAREFULLY BEFORE USING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS

May cause mild skin irritation. Moxidectin may possibly affect development and/or reproduction, and may have effects on or via lactation. Do not breathe mist, vapours or spray. Avoid skin contact. Avoid contact during pregnancy/while nursing. Handle with care. The use of chemically resistant gloves is recommended during use. Wear protective gloves and use personal protective equipment as required. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace and should be washed before reuse. Store locked up.

FIRST AID

For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water.

ENVIRONMENTAL PROTECTION

Very toxic to the aquatic environment. Avoid contamination of any water supply with product or empty container. Harmful to the soil environment. Toxic to terrestrial invertebrates. Avoid release to the environment.

DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate recycle. Do not use container for any other purpose.

EMERGENCY RESPONSE

In case of spillage wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.

EFFICACY SPECTRUM

MOXIMAX POUR ON contains moxidectin, a member of the milbemycin/avermectin family. It is effective against sensitive strains of the following internal parasites:

Cattle

Gastrointestinal Roundworms

Adult and immature forms unless otherwise stated; *Haemonchus contortus* (barber's pole worm), *Ostertagia ostertagi* (*O. lyrata*) (small brown stomach worm including inhibited larvae), *Trichostrongylus axei* (stomach hair worm), *Trichostrongylus colubriformis* (black scour worm), *Trichostrongylus longispicularis*, *Cooperia oncophora* (small intestinal worm), *Cooperia punctata*, *Oesophagostomum radiatum* (nodule worm), *Bunostomum phlebotomum* (hookworm).

Lungworm: Adult and immature *Dictyocaulus viviparus*

Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*, *Bovicola bovis*

Mange mites: *Chorioptes bovis*

Deer

Lungworm: *Dictyocaulus elerti* (*viviparus*)

Gastrointestinal roundworms: *Haemonchus contortus*, *Ostertagia*-like species*, *Trichostrongylus* spp.,

Oesophagostomum venulosum, *Chabertia ovina*

**Ostertagia leptospicularis* and *Ostertagia*-like species including *Spiculoptera* *asymetrica*, *Spiculoptera* *spiculoptera* and *Skjabinagia kalthida*.

NOTE: Resistance may develop to any anthelmintic. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance.

PERSISTENT ACTIVITY

MOXIMAX POUR ON, following a single application at the recommended dose rate, prevents reinfection of: CATTLE with *Dictyocaulus viviparus*, *Oesophagostomum radiatum* and *Bunostomum phlebotomum* for 42 days, *Ostertagia ostertagi* (*O. lyrata*) for 35 days and *Haemonchus* spp and *Trichostrongylus axei* for 28 days, and DEER with *Dictyocaulus elerti* (*viviparus*), *Ostertagia*-like spp*, *Oesophagostomum venulosum* and *Chabertia ovina* for 42 days.

**Ostertagia leptospicularis* and *Ostertagia*-like species including *Spiculoptera* *asymetrica*, *Spiculoptera* *spiculoptera* and *Skjabinagia kalthida*.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary to avoid unnecessary pain and distress when using the product other than as directed on the label.

Ready for use with standard pour-on equipment. **MOXIMAX POUR ON** should be applied along the top of the back from the shoulders to the base of the tail. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment either with scales or weighband. Choose dose according to the weight of the heaviest animals. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.

DOSEAGE AND ADMINISTRATION

Dose rate: 1mL/10kg to give 0.5mg moxidectin per kg liveweight.

BATCH:

EXP:

DOSE VOLUME TABLE

| Liveweight (kg) | Dose volume (mL) | Doses per 5 L Pack | Liveweight (kg) | Dose volume (mL) | Doses per 5 L Pack |
|-----------------|------------------|--------------------|-----------------|------------------|--------------------|
| 50 | 5 | 1000 | 250 | 25 | 200 |
| 60 | 6 | 833 | 300 | 30 | 166 |
| 70 | 7 | 714 | 350 | 35 | 142 |
| 80 | 8 | 625 | 400 | 40 | 125 |
| 90 | 9 | 555 | 450 | 45 | 111 |
| 100 | 10 | 500 | 500 | 50 | 100 |
| 110 | 11 | 454 | 550 | 55 | 90 |
| 120 | 12 | 416 | 600 | 60 | 83 |
| 130 | 13 | 384 | 650 | 65 | 76 |
| 140 | 14 | 357 | 700 | 70 | 71 |
| 150 | 15 | 333 | 750 | 75 | 66 |
| 200 | 20 | 250 | | | |

Cattle heavier than 750kg should be dosed at 1mL per 10kg liveweight.

SAFETY TO TARGET ANIMALS

MOXIMAX POUR ON has a wide margin of safety when used at the recommended dose rate.

RAINFALL

The efficacy of **MOXIMAX POUR ON** is not adversely affected if applied when the hide is wet, or if rain occurs shortly after the application. However treatment of animals under these circumstances is not recommended practice.

NON-IRRITANT

MOXIMAX POUR ON does not cause irritation to the skin or damage to the hide when used as directed.

CONCURRENT USE

MOXIMAX POUR ON is safe to use concurrently with other treatments in cattle including mineral supplements, luke treatments and vaccines. **It should not be applied at the same time as any other pour-on product.**

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MEAT: Cattle : Nil Deer: Nil Bobby calves: Nil

MILK: Nil

EPA approval number HSR001850. See www.epa.govt.nz for approval conditions.

Registered pursuant to the ACVM Act 1997, No.A011326. See www.foodsafety.govt.nz for registration conditions.

STORAGE: Store below 25°C and protect from light.

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Contains Moxidectin)

UN Number 3082, Pack Group III, Hazchem 3Z.

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call **0800 CHEMICAL (0800 243 622)**. If additional safety information is required, refer to the Safety Data Sheet. For a copy of the SDS phone 0800 100 123.

Agrecovery is available on plastic containers up to 60L and for plastic or steel large drums from 6L - 1000L.

 **AGRECOVERY**
www.agrecovery.co.nz

PHONE **0800 100 123**

