

WARNING – KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



ravensdown

Combo™ Sheep

For the treatment and control in sheep and lambs of roundworms, including benzimidazole or levamisole resistant strains. Also controls lungworms, tapeworms and liver fluke. Reduces the output of worm and fluke eggs

ACTIVE INGREDIENTS:

24.0g/litre albendazole, 37.5g/litre levamisole hydrochloride, 0.25g/litre cobalt, 2.1g/litre copper, 0.55g/litre zinc, 0.5g/litre selenium.



408603 V01



Registrant: Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz

NET CONTENTS
20 Litres

COMBO™ SHEEP READ LABEL CAREFULLY BEFORE USING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS

May be harmful if swallowed, inhaled or absorbed through the skin. May cause skin allergy. Suspected of causing genetic defects and may damage fertility or the unborn child. May cause damage to the blood and the haematopoietic system and other organ systems. Avoid contact with the skin and avoid breathing vapours. Wear protective gloves and use personal protective equipment as required. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace and should be washed before reuse.

FIRST AID

If skin contact occurs remove contaminated clothing and wash with plenty of soap and running water. For advice, or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. Harmful to the soil environment. Avoid release to the environment.

DISPOSAL

Preferably dispose of product by use. Otherwise treat the product so that it is no longer toxic and dispose of product and packaging at an approved landfill or other approved facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate recycle.

EMERGENCY RESPONSE

In case of spillage wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely.

EFFICACY SPECTRUM

COMBO SHEEP is intended for use against benzimidazole and levamisole sensitive internal parasites of the following species:
ROUNDWORMS (mature and immature): Barber's pole worm (*Haemonchus contortus*), Small brown stomach worm (*Ostertagia* spp.), Stomach hair worm (*Trichostrongylus axei*), Black scour worm (*Trichostrongylus* spp.), Small intestinal worm (*Cooperia* spp.), Hookworm (*Bunostomum* spp.), Nodule worm (*Oesophagostomum columbianum*), Thin-necked intestinal worm (*Nematodirus* spp.), Large bowel worm (*Oesophagostomum venulosum*), Large mouthed bowel worm (*Chabertia ovina*).
LUNGWORMS: Large lungworm (*Dictyocaulus filaria*)
TAPEWORMS: Heads and Segments (*Moniezia* spp.)
LIVER FLUKE (Adults): *Fasciola hepatica*
OVICIDAL: Reduces the output of viable worm and fluke eggs.
NOTE: COMBO SHEEP contains both albendazole, a member of the benzimidazole family, and levamisole. It is effective against internal parasites sensitive to one or both of these families. Resistance can develop to any anthelmintic. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce the potential for resistance to develop. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

DIRECTIONS FOR USE

By law, the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain or distress when using the product other than as directed on the label. Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do not exceed the stated dose, or dose more frequently than 3-weekly. In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species. Doses of 3 or more times those recommended can cause symptoms of Levamisole toxicity, so estimate liveweights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between Levamisole and Organophosphate dips.
 Shake container well before use. Administer orally, and check accuracy of drenching equipment before and during use. Clean drenching equipment thoroughly before and after use. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Do not under dose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. Correct drenching technique should be used.

BATCH:

EXP:

Agrecovery is available on plastic containers up to 60L and for plastic or steel large drums from 61L - 1000L.

PHONE **0800 100 123**



DOSE RATE

Sheep and Lambs: **1ml per 5kg live-weight** (4.75mg/kg albendazole, 7.5mg/kg levamisole hydrochloride).

DOSE VOLUME TABLE

Live-weight (kg)	Dose (ml)	20L pack treats (head)
16-20	4	5,000
21-25	5	4,000
26-30	6	3,333
31-35	7	2,857
36-40	8	2,500
41-45	9	2,222
46-50	10	2,000
51-55	11	1,818
56-60	12	1,666
61-65	13	1,538
66-70	14	1,428
71-75	15	1,333

Sheep heavier than 75kg should be dosed at 1ml per 5kg live-weight.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.


MEAT: Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 10 days of cessation of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 10 days following the last treatment.

EPA approval number HSRO01938. See www.epa.govt.nz for approval conditions. Registered pursuant to the ACVM Act 1997, No.A9444. See www.foodsafety.govt.nz for registration conditions.

Store locked up in the original container, tightly closed in a safe place. Store below 30°C away from direct sunlight.

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call **0800 CHEMCALL (0800 243 622)**. If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.



ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.

(Contains Albendazole)

UN Number 3082, Pack Group III, Hazchem 3Z