WARNING – KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY




Stock CopperTM

A ready-to-use liquid copper supplement for the prevention and treatment of copper deficiency in livestock

ACTIVE INGREDIENT:

50mg/ml (5%) elemental copper as copper sulphate pentahydrate





Marketed by: Ravensdown Ltd

292 Main South Road Hornby, Christchurch 8042 New Zealand Phone: 03 353 4600

ravensdown.co.nz

NET CONTENTS **20 Litres**

ravensdown **G STOCK COPPER**

STOCK COPPER[™] | READ LABEL CAREFULLY BEFORE USING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS

May be harmful if swallowed, inhaled or in contact with the skin. Causes skin irritation and may cause an allergic skin reaction. Causes serious eye irritation and may cause damage to the kidneys. Avoid contact with the skin and eyes, and do not breathe vapours. Wear protective clothing and gloves and use personal protective equipment as required. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace and should be washed before re-use.

FIRST AID

If skin contact occurs remove contaminated clothing and wash with plenty of soap and running water. If swallowed rinse mouth but do not induce vomiting. If splashed in eyes rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If inhaled remove to fresh air and keep at rest in a position comfortable for breathing. For advice, or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

ENVIRONMENTAL PROTECTION

Toxic to aquatic life with long lasting effects. Avoid release to the environment. Collect any spillage.

DISPOSAL

Preferably dispose of product by use. Otherwise treat the product so that it is no longer toxic and dispose of product and packaging at an approved landfill or other approved facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate recycle.

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Contains Copper Sulphate Pentahydrate)

UN Number 3082, Pack Group III, Hazchem 3Z EPA approval number HSR006602. See www.epa.govt.nz for approval conditions.

BATCH:

D.O.M:

Store in the original container, tightly closed, locked up, away from direct sunlight and below 30°C.

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call **0800 CHEMCALL (0800 243 622)**. If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.

DIRECTIONS FOR USE

Water treatment: Administer to adult cattle through the water supply at a rate not exceeding 5ml/head/day (250mg elemental copper), adult sheep 0.1-0.3ml/head/day (5-15mg elemental copper) and adult deer 1-2 ml/head/day (50--100mg elemental copper), unless advised by your veterinarian.

Oral drenching: For dairy cows that are drenched daily during milking, use the rates described as for water treatment above. For other stock classes, consult your veterinarian for recommended dose rates.

Addition to supplementary feed: Mix thoroughly into supplementary feed daily to supply the same rates as for water treatment above.


Foliar application: Spray pasture to be grazed with 10L Stock Copper per hectare, ideally 1-2 weeks pre-grazing, for up to 40 days supplementation. Stock Copper can also be sprayed at 10L/ha onto pasture that is due to be cut, or has been cut, for hay, silage or baleage.

Please note: Required dose rates for copper depend on several factors including dietary molybdenum, iron and zinc levels.

WARNING

Only use where copper deficiency has been diagnosed and do not use at the same time as any other copper supplements without consulting a veterinarian. Contact your veterinarian for more information and do not exceed stated dose and frequency. Frequent monitoring of your stock's copper status by liver testing is recommended.

During the facial eczema season, do not use this product at the same time as high doses of zinc as copper may interfere with the effectiveness of facial eczema control.



Agrecovery is available on plastic containers up to 60L and for plastic or steel large drums from 61L - 1000L.



PHONE **0800 100 123**