

WARNING – KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY


ravensdown

Stock IodineTM

A ready-to-use liquid iodine supplement for the prevention and treatment of iodine deficiency in livestock

ACTIVE INGREDIENT:

50mg/ml (5%) elemental iodine as potassium iodide



Marketed by: Ravensdown Ltd
292 Main South Road
Hornby, Christchurch 8042
New Zealand
Phone: 03 353 4600
ravensdown.co.nz

NET CONTENTS
20 Litres

ravensdown 

STOCK IODINE™ | READ LABEL CAREFULLY BEFORE USING

Read label carefully and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS

May cause an allergic skin reaction. Avoid contact with the skin and avoid breathing vapours. Wear protective clothing and gloves and use personal protective equipment as required. Contaminated work clothing should not be allowed out of the workplace and should be washed before re-use.

FIRST AID

If skin contact occurs remove contaminated clothing and wash with plenty of soap and running water. For advice, or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

ENVIRONMENTAL PROTECTION

Harmful to aquatic life with long lasting effects. Avoid release to the environment. Collect any spillage.

DISPOSAL

Preferably dispose of product by use. Otherwise treat the product so that it is no longer toxic and dispose of product and packaging at an approved landfill or other approved facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate recycle.

DIRECTIONS FOR USE

Water treatment: Administer to adult cattle through the water supply at a rate of 0.1-0.8ml per head/day. Adult sheep require 10-40ml per 1000 sheep/day. The higher rates are required when animals are being fed on Brassica crops.

Oral drenching: Drench ewes with 4-5ml/ewe 8 weeks and again 4 weeks pre-tup and pre-lamb. Drench other livestock at a dose rate recommended by your veterinarian. For dairy cows that are drenched daily during milking, use the rates described as for water treatment above.

Addition to supplementary feed: Mix thoroughly into supplementary feed to supply adult cattle with 0.1-0.8ml per head/day, and adult sheep with 10-40ml per 1000 sheep/day, unless advised by your veterinarian. The higher rates are required when animals are being fed on brassica crops.

Foliar application: Spray pasture to be grazed or conserved with up to 1.25L Stock Iodine per hectare, ideally 1-2 weeks pre-grazing or pre harvesting. For brassica crops that are to be grazed, spray with up to 2.5L Stock Iodine per hectare.

WARNING

Only use where iodine deficiency has been diagnosed or a need for supplementary iodine has been identified. Do not use at the same time as any other iodine supplements without consulting a veterinarian. Contact your veterinarian for more information and do not exceed stated dose and frequency. Frequent monitoring of your stock's iodine status is recommended.

EPA approval number HSR006638.
See www.epa.govt.nz for approval conditions.

BATCH:

D.O.M:

Store in the original container, tightly closed, locked up, away from direct sunlight and below 30°C.

In transport emergency call 111, Police or Fire Brigade. For 24hr specialist advice in an emergency only call **0800 CHEMCALL (0800 243 622)**. If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone 0800 100 123.

Agrecovery is available on plastic containers up to 60L and for plastic or steel large drums from 61L - 1000L.



PHONE 0800 100 123